

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



B4

(11)

EP 0 911 415 A2 = 13723

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

28.04.1999 Bulletin 1999/17

(51) Int. Cl.⁶: C12P 17/04

(21) Application number: 98121547.8

(22) Date of filing: 05.09.1991

(84) Designated Contracting States:
AT BE CH DE DK FR GB IT LI NL

(30) Priority: 17.09.1990 EP 90117842

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
91114996.1 / 0 476 442

(71) Applicant:
F. Hoffmann-La Roche AG
4002 Basel (CH)

(72) Inventors:
• Hoshino, Tatsuo
Kamakura-shi, Kanagawa-ken (JP)
• Matzinger, Peter Karl
4118 Rodersdorf (CH)

• Ojima, Setsuko
Fujisawa-shi, Kanagawa-ken (JP)
• Sugisawa, Teruhide
Yokohama-shi, Kanagawa-ken (JP)

(74) Representative:
Wibbelmann, Jobst, Dr., Dipl.-Chem.
Wuesthoff & Wuesthoff,
Patent- und Rechtsanwälte,
Schweigerstrasse 2
81541 München (DE)

Remarks:

This application was filed on 17 - 11 - 1998 as a
divisional application to the application mentioned
under INID code 62.

(54) Microbiological manufacture of L-ascorbic acid

(57) The present invention concerns a novel pro-
cess for the microbiological manufacture of L-ascorbic
acid from L-gulonolactone.


EP 0 911 415 A2

Description

[0001] The present invention concerns a novel process for the microbiological manufacture of L-ascorbic acid from L-gulono-gamma-lactone.

[0002] As pointed out above, the novel L-gulono-gamma-lactone dehydrogenase (hereinafter referred to as GLDH) provided by the present invention catalyzes the oxidation of L-gulono-gamma-lactone to L-ascorbic acid.

[0003] Enzymes which catalyze the oxidation of L-gulono-gamma-lactone (I) to L-ascorbic acid (II)



are known. Nishikimi et al. isolated L-gulono-gamma-lactone oxidase from rat liver (Arch. Biochem. Biophys., 175, 427-435, 1976), goat liver (Arch. Biochem. Biophys., 175, 427-435, 1976) and chicken kidney (Biochemistry, 21, 5076-5082, 1982). These enzymes use molecular oxygen as a direct electron acceptor in the oxidation of I to II, but the GLDH provided by the present invention does not. Moreover, any of them consists of one subunit, while the GLDH provided by the present invention consists of three kinds of subunits. Nishikimi et al. also isolated L-galactono-gamma-lactone oxidase from baker's yeast (Arch. Biochem. Biophys., 191, 479-486, 1978). This enzyme catalyzes the oxidation of both L-galactono-gamma-lactone and L-gulono-gamma-lactone to L-ascorbic acid. On the other hand, Bleeg et al. isolated L-galactono-gamma-lactone oxidase from *Saccharomyces cerevisiae* (Eur. J. Biochem., 127, 391-396, 1982). This enzyme was reported to be active on L-galactono-gamma-lactone, but not active on L-gulono-gamma-lactone. But the GLDH provided by the present invention cannot use L-galactono-gamma-lactone as its substrate.

[0004] Thus, the GLDH provided by the present invention is clearly distinct from L-galactono-gamma-lactone oxidase of yeast in its substrate specificity. On the other hand, Shigeoka et al. reported the characteristics of crude L-gulono-gamma-lactone dehydrogenase of *Euglena gracilis* z (Agric. Biol. Chem., 43, 2187-2188, 1979), i.e. the enzyme catalyzed the oxidation of both L-gulono-gamma-lactone and L-galactone-gamma-lactone, and was incapable of using oxygen as an electron acceptor. Furthermore, it is common knowledge that algae are difficult to handle due to the problems encountered with the growth of these microorganisms (fragility, cell division and the extended time periods involved).

[0005] The novel GLDH provided by the present invention clearly differs from it in its specificities of substrate. In addition, there exists no report on the isolation of L-gulono-gamma-lactone dehydrogenase of *Euglena gracilis* z up to now.

[0006] There have been no-reports up to now on the conversion of L-gulono-gamma-lactone to L-ascorbic acid by using bacteria. Accordingly to the present invention, however, it has been found that bacteria are capable of producing L-ascorbic acid from L-gulono-gamma-lactone. This is the first possibility of such production of L-ascorbic acid from L-gulono-gamma-lactone by using bacteria.


[0007] As described above, no disclosure exists of a purified enzyme, e.g. of bacterial origin, which has dehydrogenase activity to oxidize L-gulono-gamma-lactone to L-ascorbic acid. It has been found that the purified enzyme, e.g. isolated from the soluble fraction of bacterial cells of specific microorganisms, catalyzes the oxidation of L-gulono-gamma-lactone to L-ascorbic acid. The present invention has been accomplished on the basis of this finding.

[0008] As pointed out, one of the objects of the present invention is to provide a process for producing L-ascorbic acid by fermentation by means of bacteria.

[0009] The physico-chemical properties of the purified novel GLDH, e.g. as exemplified below are as follows:

1) Enzyme activity:

The novel GLDH of the present invention catalyzes the oxidation of L-gulono-gamma-lactone to L-ascorbic acid in the presence of an electron acceptor according to the following reactions:



The enzyme does not utilize oxygen as an electron acceptor. This was affirmed by the missing catalyzing activity of the enzyme to convert L-gulono-gamma-lactone to L-ascorbic acid using oxygen as said electron acceptor. Furthermore, no oxygen consumption was detected in the reaction mixture as detected by an oxygen probe. However, any conventional compound which has the ability to act as an electron acceptor can be utilized in conjunction with the enzyme of this invention. As such electron acceptor, a coenzyme, or a compound exhibiting such coenzyme function, e.g. 2,6-dichlorophenolindophenol (hereinafter referred to as DCIP), phenazine methosulphate, Wurster's blue, ferricyanide, coenzyme Q, or cytochrome c, etc. can be used.

The enzyme assay was performed at 25 °C by measuring the decrease of absorbance at 600 nm of DCIP spectrophotometrically. One unit of enzyme activity was defined as the amount of enzyme which catalyzed the reduction of 1 µmole of DCIP per minute. The extinction coefficient of DCIP at pH 7.0 was taken as 14.5 mM⁻¹. A cuvette with 1 cm light path contained 0.16 mM DCIP, 1.6 mM phenazine methosulfate, 200 mM potassium phosphate buffer, 400 mM L-gulono-gamma-lactone, enzyme solution and water in a final volume of 0.5 ml. A reference cuvette contained all components except L-gulono-gamma-lactone. The reaction was started by the addition of L-gulono-gamma-lactone. Enzyme activity was measured as the initial reduction rate of DCIP.

2) Substrate specificity:

Substrate-specificity of the enzyme was determined using the same enzyme assay method as described as above except that the various substrate solutions (400 mM) were used instead of L-gulono-gamma-lactone. The results of the measurement are shown in Table 1. The GLDH is highly active on L-gulono-gamma-lactone and D-xylose and weakly active on D-glucurono-gamma-lactone, D-glucose and D-mannose.

3) Optimum pH:

The correlation between the reaction rate of the GLDH and pH was determined using the same enzyme assay method as described under 1) except that various pH's buffers were used. The results are shown in Table 2. The enzyme showed the highest enzyme activity in a pH range between 7.0 and 8.0.

4) pH stability:

The purified enzyme was kept standing in buffers of various pH for 192 hours at 4 °C. The residual activity was measured using the same enzyme assay method as described under 1). The results of the measurement are shown in Table 3. The purified enzyme was relatively stable at any pH between 6.5 and 9.2.

5) Heat stability:

The purified enzyme was treated for 5 minutes at various temperatures in 200 mM potassium phosphate buffer (pH 7.0), and then cooled immediately in ice water. The residual activity was measured using the same enzyme assay method as described under 1). The results are shown in Table 4. The purified enzyme was stable up to 30 °C, and lost about 50 and 80% of its activity after incubation at 55° and 60°C, respectively.

6) Optimum temperature:

The enzyme activities of GLDH were measured at temperatures from 25 to 55 °C by the same enzyme assay method as described under 1). The results are shown in Table 5. The enzyme did not have a distinct optimum temperature in the range tested.

7) Molecular weight:

The molecular weight of the GLDH was determined by high performance liquid chromatography using a size exclusion gel column (TSK gel G3000 SWxL column, 7.8 mm x 30 cm) equilibrated with 100 mM potassium phosphate buffer (pH 7.0) containing 300 mM sodium chloride. As molecular weight standards, cyanocobalamin (M.W. 1,350), myoglobin (M.W. 17,000), ovalbumin (M.W. 44,000), gamma-globulin (M.W. 158,000) and thyroglobulin (M.W. 670,000) were used. The molecular weight of the GLDH was determined to be 110,000 ± 2000.

Next, subunit-components of the purified GLDH were determined. The purified GLDH was treated by sodium dodecyl sulfate (hereinafter referred to as SDS) in the presence of beta-mercaptoethanol, and applied on the same column as described above equilibrated with 100 mM sodium phosphate buffer (pH 7.0) containing 0.1 % SDS. As molecular weight standards, lysozyme (M.W. 14,000), soy bean trypsin inhibitor (M.W. 21,500), carbonic anhydrase (M.W. 31,000), ovalbumin (M.W. 45,000), bovine serum albumin (M.W. 66,200) and phosphorylase B (M.W. 92,500) were used. The enzyme consists of three subunits whose molecular weights were 61,000, 32,500 and 16,500. The sum of these molecular weights is 110,000, the total molecular weights of native GLDH.

The largest component (M.W. 61,000) is probably a flavoprotein, since upon gel electrophoresis in SDS it showed intense fluorescence when the unstained gel was exposed to ultraviolet light. The second component (M.W. 32,500), which was stained by heme staining, is a cytochrome.

The third compound (M.W. $16,500 \pm 500$) is a simple protein, i.e. a protein carrying no prosthetic group. To sum up, the respective molecular weights are

61,000 \pm 1,000
 32,500 \pm 1,000
 16,500 \pm 500,

the standard deviations having been established by conventional means, e.g. SDS electrophoresis.

The heme staining of the above electrophoresis gel was carried out according to the procedure described by P.E. Thomas et al. in *Analytical Biochemistry* 75, 168-176 (1976), which can be summarized as follows:

A 6.3 mM 3,3',5,5'-tetramethylbenzidine (TMBZ) solution was freshly prepared in methanol. Immediately before use, 3 parts of the TMBZ solution were mixed with 7 parts of 0.25 M sodium acetate, pH 5.0. The gel was immersed into this mixture at room temperature in the dark for 2 hours with occasional mixing. H_2O_2 was added to a final concentration of 30 mM to stain the second protein component including a cytochrome.

8) Absorption spectrum:

The absorption spectrum of the purified GLDH reduced with sodium dithionite showed maxima at 416, 521 and 552 nm in the visible region, indicating the presence of a cytochrome c component, as shown in Fig. 1.

9) Measurement of the K_m value:

Using the same enzyme assay method as described above, the velocity of the oxidizing reaction with varying the concentrations of L-gulono-gamma-lactone from 0.18 mM to 90 mM was measured to determine the K_m value for L-gulono-gamma-lactone. The maximum reaction velocity was found at the substrate concentration of about 71.8 mM. The apparent Michaelis constant (K_m) was calculated to be 34.8 mM with DCIP as the electron acceptor.

10) Effect of metal ions:

Using the same enzyme assay method as described under 1), the effect of various metal ions on the enzyme activity was examined. The results are shown in Table 6. Cu^{2+} and Mn^{2+} showed strong inhibition of the enzyme.

11) Effect of inhibitors:

Using the same enzyme assay method as described above, the effect of various inhibitors on the enzyme activity was examined. The results are shown in Table 7. No compounds tested gave an inhibitory effect on the GLDH.

12) Purification method:

Purification of L-gulono-gamma-lactone dehydrogenase is effected by the combination of known purification methods, such as ion exchange chromatography, liquid chromatography, adsorption chromatography, gel-filtration chromatography, gel-electrophoresis, salting out and dialysis.

[0010] The microorganism used include all strains belonging to the genus *Gluconobacter*, showing good growth when cocultured in the presence of *Bacillus megaterium*. Mutants and variants of said microorganism can be also used in the present invention. The preferred strain is *Gluconobacter oxydans*.

[0011] The strains have been denominated and classified as *Gluconobacter oxydans* by reference to *Bergey's Manual of Determinative Bacteriology*, 8th edition, 1974, and, in particular, in view of the fact that they exhibit the following characteristics:

- a) 2-keto-L-gulonic acid is produced from L-sorbose,
- b) ethanol is oxidized to acetic acid,
- c) D-glucose is oxidized to D-gluconic acid and 2-keto-D-gluconic acid,
- d) ketogenesis of polyalcohols,
- e) pellicle and ring growth in mannitol broth (24 hours cultivation) at pH 4 and 5, and pellicle growth in D-glucose broth at pH 4.5.

In addition, they exhibit the following properties:

- f) dihydroxyacetone is not substantially produced from glycerol,
- g) 2-keto-D-glucaric acid is produced from D-sorbitol and D-glucaric acid, but not from D-glucose, D-fructose, D-gluconic acid, D-mannitol or 2-keto-D-gluconic acid,
- h) polymorphic, no flagella observed,
- i) a brown pigment is produced from D-fructose,
- j) good growth when co-cultured in the presence of *Bacillus megaterium* or a cell extract thereof.

k) streptomycin sensitive.

[0012] - The specific and preferred *Gluconobacter oxydans* strain has been deposited at the Deutsche Sammlung von Mikroorganismen in Goettingen under DSM 4025 on March 17, 1987.

[0013] The cells of the *Gluconobacter oxydans* strain are rodshaped with round ends. The diameter of a cell of the *Gluconobacter oxydans* strain is, on the average, about 0.3 - 0.6 μm , its length about 0.9 - 1.6 μm , mainly 1 - 1.5 μm .

[0014] For the preparation of the GLDH, the microorganism may be cultured in an aqueous medium supplemented with appropriate nutrients under aerobic condition. The cultivation may be conducted at a pH between about 4.0 and 9.0, preferably between about 6.0 and 8.0. While cultivation period varies depending upon pH, temperature and nutrient medium used, usually 2 to 5 days will bring about favorable results. A preferred temperature range for carrying out for the cultivation is from about 13 to 36°C, preferably from 18 to 33°C.

[0015] It is usually required that the culture medium contains such nutrients as assimilable carbon sources, digestible nitrogen sources and inorganic substances, vitamins, trace elements and other growth promoting factors. As assimilable carbon sources, L-sorbose, glycerol, D-glucose, D-mannitol, D-fructose, D-arabitol and the like can be used. Various organic or inorganic substances may also be used as nitrogen sources, such as yeast extract, meat extract, peptone, casein, corn steep liquor, urea, amino acids, nitrates, ammonium salts and the like. As inorganic substances, magnesium sulfate, potassium phosphate, ferrous and ferric chlorides, calcium carbonate and the like may be used.

[0016] In the following, an embodiment for isolation and purification of the GLDH from the above specified microorganisms after the cultivation is briefly described.

(1) Cells are harvested from the fermentation broth by centrifugation or filtration.

(2) The cells are suspended in the buffer solution and disrupted by means of a homogenizer, sonicator or treatment with lysozyme and the like, to give a solution of disrupted cells.

(3) The GLDH is isolated and purified from cell free extract of disrupted cells, preferably from the soluble fraction of microorganisms.

[0017] In these steps, one preferably makes use of column chromatography, e.g.

- 1) DEAE cellulose column chromatography,
- 2) Q-Sepharose column chromatography,
- 3) Hydroxylapatite column chromatography,
- 4) Sephacryl S-300 column chromatography,
- 5) polyacrylamide gel electrophoresis, etc.

[0018] The GLDH provided by the present invention is used as a catalyst for the production of L-ascorbic acid from L-gulonogamma-lactone. This reaction should be conducted at pH values of from about 6.0 to 9.0 in the presence of an electron acceptor, for example, DCIP, PMS, Wurster's blue, ferricyanide, coenzyme Q, cytochrome c and the like, in a solvent such as McIlvaine buffer, potassium phosphate buffer, Tris-HCl buffer and the like.

[0019] A preferred temperature range for carrying out the reaction is from about 25 to 55 °C. When the pH and the temperature are set at about 7.0 - 8.0 and 30 - 50°C, respectively, the reaction usually brings about the most preferable results. Concentration of L-gulonogamma-lactone as substrate in a solvent varies depending on other reaction conditions, but, in general, is desirable to be about 10 - 150 g/l, most preferably from about 10 - 100 g/l.

[0020] For this reaction, the enzyme may also be used in an immobilized state with an appropriate carrier. Any means of immobilizing enzyme generally known to the art may be used. For instance, the enzyme may be bound directly to a membrane, granules or the like of a resin having functional group(s), or it may be bound through bridging compounds having bifunctional group(s), for example, glutaraldehyde, to the resin.

[0021] In as far as the fermentation process is concerned, the following parameters apply and the process is suitably done according to the following procedure respectively: a microorganism with the capability to produce II is cultivated in a nutrient aqueous solution in presence of compound I or is, after its growth, brought into contact with I in a buffer solution and then further incubated.

[0022] Examples of microorganisms are: e.g.

Bacteria, e.g. of the genus:

Acetobacter, e.g. *Acetobacter suboxydans* (DSM 5935) [date of deposit: May 5, 1990], *Acetobacter oxydans* (DSM 5936) [May 5, 1990], *Acetobacter melanogenus* (NCIMB 8086)

Gluconobacter, e.g. *Gluconobacter oxydans* (ATCC 621).

Gluconobacter oxydans (DSM 4025).

[0023] Further suitable bacteria are: e.g.

Actinomyces, e.g. of the genus Streptomyces, such as Streptomyces antibioticus (ATCC 8633), Streptomyces eurocidicus (ATCC 19551),

Streptomyces lavendulae (DSM 5926) [May 5, 1990], Streptomyces olivaceus (ATCC 3335), Streptomyces netropsis (NRRL 2268).

[NRRL = Northern Utilization Research and Development Division of U.S.D.A., Peoria, Illinois, USA

ATCC = American Type Culture Collection, Rockville, Maryland, USA

NCIMB = National Collection of Industrial + Marine Bacteria, Torry Research Station, Aberdeen AB 9 8DG, Scotland

DSM = German Collection of Microorganisms and Cell Cultures, Braunschweig]

[0024] Specially selected for the reaction were species of the genus Acetobacter and Gluconobacter, especially the strains Acetobacter suboxydans and Gluconobacter oxydans. These strains have been deposited for the safeguard of the process.

[0025] It is understood that every microorganism used according to the invention should preferably be grown in a nutrient medium before it is used for the inventory fermentation process. Its growth is possible in aqueous medium.

[0026] The nutrient medium with the grown microorganism can be used directly for the fermentation reaction.

[0027] The composition of the inventive reaction medium can be much simpler, e.g. a solution of educt I combined with the separately grown microorganisms in a buffer solution without any further additions.

[0028] The pH of the medium should preferably be between 2 and 9, preferably ca. 4 - 7. If desired the pH value may be adjusted by a buffer system.

[0029] For optimal yields it is preferable to use the educt in a concentration between ca. 1 and ca. 10 % (per weight).

[0030] The preferable fermentation time is between 2 and 100 hours, especially between 4 and 72 hours. A feeding of educt I may prolong the fermentation time.

[0031] It is understood that the oxidation represents an aerobic process. These conditions are fulfilled with vivid shaking or stirring of the reaction medium under air or under additional aeration.

[0032] The following examples illustrate the present invention.

Example 1

Preparation of L-gulono-gamma-lactone dehydrogenase

(1) Cultivation of Gluconobacter oxydans DSM 4025

[0033] Gluconobacter oxydans DSM 4025 was grown on an agar slant medium containing mannitol 5.0 %, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.25 %, corn steep liquor 1.75 %, baker's yeast 5.0 %, urea 0.5 %, CaCO_3 0.5 % and agar 2.0 % at 27°C for 4 days. One loopful of the agar slant culture of Gluconobacter oxydans DSM No. 4025 was inoculated into 50 ml of a seed culture medium containing L-sorbose 8.0 %, glycerol 0.05 %, urea 0.5 %, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.25 %, corn steep liquor 1.75 %, baker's yeast 5.0 % and CaCO_3 1.5 % in a 500 ml-Erlenmeyer flask, and cultivated at 30°C for 1 day on a rotary shaker (180 rpm). Five ml of this culture were transferred into 50 ml of the same medium in a 500 ml-Erlenmeyer flask, and cultivated in the same manner as described above. Two liters of the seed culture thus prepared were used as an inoculum for a 30 l-jar fermentor containing 20 liters of medium containing L-sorbose 8.0 %, glycerol 0.05 %, urea 1.2 %, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.25 %, corn steep liquor 3.0 %, baker's yeast 5.0 % and CaCO_3 1.5 %. The jar fermentor was operated at 30°C, with 400 rpm agitation and 0.5 vvm (volume of air/ volume of medium/minute) for aeration. After 40 hours fermentation, the cultured broth was centrifuged at 1,500 rpm for 10 minutes to remove calcium carbonate, then at 8,000 rpm (10,000 x g) to pellet the cells. The cell cake was washed with 0.85 % NaCl solution once. From 20 liters of the broth, about 100 g (wet weight) of the cells were obtained.

(2) Preparation of the soluble fraction

[0034] The cells of Gluconobacter oxydans DSM 4025 (95 g, wet weight) from the above step (1) were washed with 0.85 % NaCl solution twice. The washed cells were suspended in 380 ml of 10 mM potassium phosphate buffer (pH 7.0), and the cell suspension was homogenized with a French press homogenizer at 1,500 kg/cm². Cell debris was removed by centrifugation at 1,800 x g for 10 minutes, and then the supernatant (hereinafter referred to as cell free extract) was centrifuged at 100,000 x g for 60 minutes. The resulting supernatant (450 ml) was collected as the soluble

fraction of the cells of *Gluconobacter oxydans* DSM 4025.

(3) Diethylaminoethyl (hereinafter referred to as DEAE)cellulose column chromatography

5 [0035] The soluble fraction (450 ml) obtained in the previous step was dialyzed against 10 mM potassium phosphate buffer (pH 7.0). The dialysate (500 ml) was applied to a DEAE-cellulose column (2.5 x 120 cm), equilibrated with 10 mM potassium phosphate buffer (pH 7.0). The column was washed with the same buffer and then with the same buffer containing 0.25 M NaCl. The GLDH was eluted with the same buffer containing 0.5 M NaCl.

10 (4) Q-Sepharose column chromatography

[0036] The pooled active fraction (200 ml) from the previous step was dialyzed against two batches of 2 liters of 10 mM potassium phosphate buffer (pH 7.0), and applied to a Q-Sepharose column (2.5 x 50 cm) equilibrated with the same buffer. After the column was washed with the same buffer, the GLDH was eluted with the linear gradient of NaCl
15 from 0.3 M to 0.5 M.

(5) Hydroxylapatite (Bio-gel HTP) column chromatography

[0037] The pooled active fraction (210 ml) from the previous step was dialyzed against two batches of 2 liters of 10 mM potassium phosphate buffer (pH 7.0), and applied on a hydroxyl apatite column (2.5 x 25 cm) equilibrated with the same buffer. The column was washed with the same buffer, and the GLDH was eluted with the linear gradient of potassium phosphate buffer (pH 7.0) from 10 mM to 40 mM. Fractions having enzyme activity were combined and concentrated to about 20 ml by ultrafiltration using a ultrafilter (pM10, Amicon).

25 (6) Sephacryl S-300 column chromatography

[0038] A portion of the enzyme fraction (2 ml) from the previous step was applied on Sephacryl S-300 column (1.0 x 100 cm) equilibrated with 50 mM potassium phosphate buffer (pH 7.0) containing 50 mM NaCl, and developed with the same buffer. The fractions containing electrophoretically homogeneous GLDH were combined and stored at -80°C.

30 [0039] Using the purification steps as described above, the GLDH was purified about 900-fold. Summary of the purification steps of GLDH is shown in Table 8.

(7) Purity of the isolated enzyme

35 [0040] To estimate the purity of the GLDH isolated, a polyacrylamide gel electrophoresis (separation gel: 10 % polyacrylamide; conditions of electrophoresis: 20 mA at 4°C for 6 hours) was performed. The enzyme yielded a single band stained by Coomassie brilliant blue R-250. The protein band showed GLDH activity when the unstained gel was immersed in 50 mM potassium phosphate buffer (pH 7.0) containing 50 mM L-gulono-gamma-lactone, 20 µg/ml of nitroblue tetrazolium and 40 µg/ml of phenazine methosulfate for 20 minutes.

40 (8) Identification of the reaction product

[0041] The reaction mixture containing 0.5 ml of the purified GLDH (7.5 µg, 469 units/mg protein), 1.0 ml of 0.5 M potassium phosphate buffer (pH 7.0), 0.1 g of L-gulono-gamma-lactone, 0.1 ml of 10 mM phenazine methosulfate, and water in a final volume of 2.0 ml was incubated for 2 hours at 30°C. The reaction product was analyzed by both thin layer chromatography and high performance liquid chromatography. Thin layer chromatography was done as follows: a sample (1 µl) was spotted on a silica gel plate (Merck, U.S.A.), developed with a solvent system of n-propanol-water-1 % phosphoric acid-formic acid (400 : 100 : 10 : 1) at room temperature for 2 hours. The plate was then dried and observed under an ultra-violet lamp. The product was found as UV-absorption spot at a R_f value of about 0.7 that corresponded
50 with an authentic sample of L-ascorbic acid. High performance liquid chromatography was done as follows: a sample was applied on a LiChrosorb NH₂ column (Merck, U.S.A., 0.4 x 25 cm) equilibrated with a solvent system of acetonitrile-water-acetic acid 87 : 11 : 2. The flow rate was set at 3.0 ml/min, and the detection of products was made at 254 nm. As a result, the product eluted at the same retention time as an authentic sample of L-ascorbic acid.

[0042] Consequently, the product was identified to be L-ascorbic acid.

55 [0043] The productivity of L-ascorbic acid was 0.71 g/hour.

Example 2

L-Ascorbic acid production from L-gulonogamma-lactone by fermentation (growing cell).

[0044] A 200 ml of the seed culture of *Gluconobacter oxydans* DSM 4025 prepared in the same manner as described in Example 1-(1) was used to inoculate 2 liters of medium, which contained L-gulono-gamma-lactone 8 %, glycerol 0.05 %, baker's yeast 5.0 %, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.25 %, corn steep liquor 1.75 %, urea 0.5 % and CaCO_3 1.5 % (the initial pH set at 7.0), in a 3-l jar fermentor. Fermentation was conducted at 30 °C, 700 rpm for agitation and 0.5 vvm for aeration. As shown in Table 9, 8.6 g/l of L-ascorbic acid was produced in 66 hours of fermentation.

Example 3

L-Ascorbic acid production from L-gulono-gamma-lactone under a resting cell system.

[0045] The cells of *Gluconobacter oxydans* DSM 4025 prepared in the same manner as described in Example 1-(1), 0.1 g to 0.67 g, were added into 50 mM potassium phosphate buffer (pH 7.0) containing 47.6 mg/ml or 89.3 mg/l of L-gulono-gamma-lactone in a total volume of 3 ml. The reaction mixture was incubated for 6 hours at 30°C with shaking (280 rpm). The results are shown in Table 10. The highest productivity of L-ascorbic acid was 20 mg/hour/g-cells. And the highest yield of L-ascorbic acid was 13.92 g/l.

Example 4

L-Ascorbic acid production from L-gulono-gamma-lactone using the cell free extract of *Gluconobacter oxydans* DSM 4025.

[0046] The reaction mixture containing 1.0 ml of cell free extract (protein content: 10.3 mg/ml) of *Gluconobacter oxydans* DSM 4025, as prepared in the same manner as described in Example 1-(1) and (2), 1 ml of 0.5 M potassium phosphate buffer (pH 7.0) and 0.5 ml of 17.8 % L-gulono-gamma-lactone was incubated at 30°C for 17.5 hours. As a result, 2.19 g/l of L-ascorbic acid was produced.

Example 5

(growing cell system).

[0047] The bacterium *Acetobacter suboxydans* (DSM 5935) is grown on an agar slant in medium 1 (composition: 50 % yeast water, 5 % mannitol, in tap water adjusted to pH 6.5). After 2 days incubation at 30°C one loopful of cells are used to inoculate 5 ml liquid medium 1 (the same composition as described above but without agar). The tube is shaken at 30°C with 220 rpm for 3 days. 1 ml preculture is used to inoculate 100 ml medium 1 together with 1 g of educt I. The flasks are shaken at 30°C with 220 rpm. The process was terminated after 72 hours. According to the analytical determinations 3 % educt I have been converted into product II.

Example 6

(resting cell system).

[0048] The bacterium *Acetobacter suboxydans* (DSM 5935) is grown as described above. The 100 ml cultures are harvested by centrifugation (10.000 rpm, 10 min.) and frozen in plastic vials above liquid nitrogen. 100 ml flasks with medium 1 are inoculated with 0.5 ml each from the vials, shaken at 30°C at 220 rpm and 1 g educt I is added at the same time. According to the analytical determinations 10 % of the educt I is transformed into product II within 72 hours.

Example 7

(resting cell system).

[0049] The bacterium *Acetobacter suboxydans* (DSM 5935) is grown as described in Example 6. The 100 ml cultures, inoculated with 0.5 ml from the vials serve as preculture for another growth culture in the same medium. These flasks with 100 ml medium 1 are inoculated with 5 ml preculture and shaken for 24 hours at 30°C with 220 rpm. The grown cells are harvested by centrifugation (10,000 rpm, 10 min.). 3 g of wet cells (corresponds to cells from one flask) are

suspended in a buffer solution (pH 6.0, 0.05M phosphate buffer) together with 0.1 g of educt I (added as powder) in a total volume of 10 ml. 30 % of educt I is transformed within 48 hours into product II.

Example 8

(resting cell system).

[0050] The bacterium *Acetobacter suboxydans* (DSM 5935) is grown on agar slants as described in Example 1. The layer of grown cells is suspended in 10 ml of a physiological NaCl solution (0.9 %). This suspension is used to inoculate 10 shake flasks (100 ml medium 1) with 1 ml each. The flasks are incubated for 4 days at 30°C with 220 rpm. The 10 cultures are used to inoculate a 10 lt blade-stirred bioreactor (9,000 ml medium 1). Growth conditions: temperature: 30°C, aeration 0.4 vvm, stirring 500 rpm. After 44 hours of growth the cells are harvested by centrifugation (continuous centrifugation with 12,000 rpm). The yield was 21 g wet cells per liter. The cells are deep frozen in portions. In the example 1 g of the frozen cells is thawed quickly and added together with 0.4 g educt I to 0.05M phosphate buffer of pH 7.0. The total volume of the assay mixture was 5 ml. According to analytical determinations 19 % of I were transformed into product II.

Table 1

Substrate Specificity of L-Gulono-r-lactone Dehydrogenase	
Substrate	Relative activity (%)
L-Gulono-r-lactone	100
L-Galactono-r-lactone	0
D-Glucurono-r-lactone	6.38
D-Glucono-δ-lactone	0
D-Glucuronic acid	0
D-Gluconic acid	0
D-Glucose	23.4
D-Mannose	7.23
D-Galactose	0
L-Gulose	0
D-Xylose	110.6

Table 2

Optimum pH of L-Gulono-r-lactone Dehydrogenase			
pH value	Relative activity (%)		
	Buffers		
Mcllvain	Potassium phosphate	Tris-HCl	
4.0	0	-	-
4.5	0	-	-
5.0	8.2	-	-
5.5	18.2	-	-
6.0	58.5	-	-
6.5	77.8	56.4	-
7.0	100	83.6	45.9
7.5	105.8	102.3	58.2
8.0	109.3	-	61.7
8.5	-	-	60.0

(- : not determined)

Table 3

pH Stability of L-Gulono-r-lactone Dehydrogenase				
pH value	Relative activity (%)			
	Buffers			
	Mcllvaine	Potassium phosphate	Tris-HCl	NH ₄ OH-NH ₄ Cl
4.0	0	-	-	-
4.5	0	-	-	-
5.0	0	-	-	-
5.5	9.38	-	-	-
6.0	48.4	62.5	-	-
6.5	75.0	50.3	-	-
7.0	79.7	71.9	84.4	-
7.5	100	95.3	75.0	-
8.0	84.4	-	62.5	-
8.3	-	-	53.2	57.8
8.85	-	-	-	115.6
9.2	-	-	-	100
9.7	-	-	-	68.8

Table 4

Temperature Stability of L-Gulono-r-lactone Dehydrogenase	
Temperature (°C)	Relative activity (%)
0	100
25	100
30	100
35	89.6
40	87.5
45	72.9
50	70.8
55	53.1
60	18.8
65	0
70	0
75	0

Table 5

Optimum Temperature of L-Gulono-r-lactone Dehydrogenase	
Temperature (°C)	Relative activity (%)
25	100
30	103.2
35	101.8
40	106.6
45	101.4
50	115.1
55	102.7

Table 6

Effect of Various Metals or L-Gulono-r-lactone Dehydrogenase					
Metal	Concentration (mM)	Relative activity (%)	Metal	Concentration (mM)	Relative activity (%)
Ca(NO ₃) ₂ · 4H ₂ O	0.19	100	MgSO ₄ · 7H ₂ O	0.19	92
	0.38	100		0.38	96
	0.89	96		0.89	92

Table 6 (continued)

Effect of Various Metals on L-Gulono-r-lactone Dehydrogenase					
Metal	Concentration (mM)	Relative activity (%)	Metal	Concentration (mM)	Relative activity (%)
CaCl ₂	0.19	96	MgCl ₂ · 6H ₂ O	0.19	96
	0.38	104		0.38	96
	0.89	96		0.89	96
CoCl ₂ · 6H ₂ O	0.19	102	MnCl ₂ · 4H ₂ O	0.19	100
	0.38	110		0.38	0
	0.89	100		0.89	0
CoSO ₄ · 7H ₂ O	0.19	96	MnSO ₄ · 4-6H ₂ O	0.19	100
	0.38	108		0.38	0
	0.89	88		0.89	0
CuSO ₄	0.19	92	Na ₂ MoO ₄ · 2H ₂ O	0.19	104
	0.38	24		0.38	96
	0.89	12		0.89	88
CuSO ₄ · 5H ₂ O	0.19	80	TiCl ₄	0.19	88
	0.38	20		0.38	92
	0.89	0		0.89	80
Cu(NO ₃) ₂ · 3H ₂ O	0.19	88	ZnCl ₂	0.19	88
	0.38	24		0.38	88
	0.89	0		0.89	80
CuCl ₂ · 2H ₂ O	0.19	84	ZnSO ₄ · 7H ₂ O	0.19	88
	0.38	8		0.38	88
	0.89	0		0.89	76
Fe(SO ₄) ₃ · xH ₂ O	0.19	104	NiSO ₄ · 7H ₂ O	0.19	96
	0.38	92		0.38	96
	0.89	76		0.89	88
K ₄ Fe(CN) ₆ · 3H ₂ O	0.10	0	None		100

Table 7

Effect of Various Inhibitors on L-Gulono-r-lactone Dehydrogenase Activity		
Inhibitor	Concentration (mM)	Relative activity (%)
EDTA	0.96	95.2
	1.89	100
	4.46	90.5

Table-7 (continued)

Effect of Various Inhibitors on L-Gulono-r-lactone Dehydrogenase Activity		
Inhibitor	Concentration (mM)	Relative activity (%)
N-Ethyl-maleimide	0.96	95.2
	1.89	100
	4.46	100
Sodium azide	0.96	100
	1.89	97.6
	4.46	97.6
Monoiodoacetate	0.96	104.8
	1.89	100
	4.46	92.9
PCMB (p-chloro-mercuribenzoate)	0.96	90.5
	1.89	102.4
	4.46	100
Na ₂ HAsO ₄ 7H ₂ O	0.96	95.2
	1.89	104.8
	4.46	97.6
Sodium fluoride	0.96	107.1
	1.89	114.3
	4.46	97.6
KCN	4.46	97.6
	0.96	96.0
	1.89	96.0
Hydroxylamine hydrochloride	4.46	96.0
	1.0	100
	2.0	100
Hydrazine monohydrate	1.0	100
None		100

Table 8

Summary of Purification Steps of L-Gulono-r-lactone Dehydrogenase				
Step	Total activity (units)	Total protein (mg)	Specific activity (units/mg)	Recovery (%)
Soluble fraction	1179.9	2289.5	0.52	100
DEAE Cellulose	766.9	66.44	11.54	65.0
Q Sepharose	761.8	8.65	88.10	64.6
Hydroxylapatite	473.1	3.24	146.0	40.1
Sephacryl S-300	37.8	0.0806	469.0	-

Table 9

L-Ascorbic Acid Production from L-Gulono-r-lactone by Fermentation	
Cultivation time (hour)	L-Ascorbic acid produced (g/l)
0	0.0715
18.5	3.54
27	5.12
42	6.96
51	7.88
66	8.57
75	8.46
90	7.72

Table 10

L-Ascorbic Acid Production from L-Gulono-r-lactone by Resting Cell System			
Initial L-gulono-r-lactone concentration (mg/ml)	Cell concentration (g cell wet weight/ml)	L-Ascorbic acid produced (mg/ml)	
		at 3 hr	at 6 hr
0	0.1	0	0
47.6	0.1	5.98	7.22
	0.17	6.46	7.82
	0.33	7.83	9.65
	0.67	9.51	11.28
89.3	0.1	5.88	7.62
	0.17	6.88	9.50
	0.33	8.37	13.92
	0.67	8.67	13.51

Claims


1. Process for the manufacture of L-ascorbic acid, which process comprises oxidizing L-gulono-gamma-lactone with the aid of bacteria, or their cell free extracts, or the soluble fraction of the cells.
2. Process according to claim 1, employing bacteria of the genera *Acetobacter*, *Gluconobacter* or *Streptomyces*, a functional equivalent, subculture, mutant, variant or cell free extract thereof.
3. Process according to claim 2, wherein *Acetobacter suboxydans* or *Gluconobacter oxydans* is used.
4. A process for producing L-ascorbic acid which comprises oxidizing L-gulono-gamma-lactone with the aid of a microorganism having the identifying characteristics of the strain *Gluconobacter oxydans* DSM 4025, a functional

equivalent, subculture, mutant, variant or cell free extract thereof, in a medium in the presence of L-gulono-gamma-lactone as a substrate, and isolating L-ascorbic acid from the cultivation medium.

- 5 5. A process according to claim 4, wherein the microorganism is cultured in a medium containing L-gluono-gamma-lactone and appropriate nutrients.
6. A process according to claim 5, wherein the concentration of the substrate is 10 - 150 g, preferably 10 to 100 g/l.
7. A process according to claim 6, wherein L-ascorbic acid is produced in a concentration of more than 3 - 14 g/l.

Fig. 1.

Absorption Spectra of L-Gulonono-r-lactone Dehydrogenase



(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 911 415 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:

06.05.1999 Bulletin 1999/18

(51) Int. Cl.⁶: **C12P 17/04**

(43) Date of publication A2:

28.04.1999 Bulletin 1999/17

(21) Application number: **98121547.8**

(22) Date of filing: **05.09.1991**

(84) Designated Contracting States:

AT BE CH DE DK FR GB IT LI NL

(30) Priority: **17.09.1990 EP 90117842**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

91114996.1 / 0 476 442

(71) Applicant:

**F. Hoffmann-La Roche AG
4002 Basel (CH)**

(72) Inventors:

• **Hoshino, Tatsuo
Kamakura-shi, Kanagawa-ken (JP)**

• **Matzinger, Peter Karl**

4118 Rodersdorf (CH)

• **Ojima, Setsuko**

Fujisawa-shi, Kanagawa-ken (JP)

• **Sugisawa, Teruhide**

Yokohama-shi, Kanagawa-ken (JP)

(74) Representative:

Wibbelmann, Jobst, Dr., Dipl.-Chem.

Wuesthoff & Wuesthoff,

Patent- und Rechtsanwälte,

Schweigerstrasse 2

81541 München (DE)

(54) **Microbiological manufacture of L-ascorbic acid**

(57) The present invention concerns a process for the microbiological manufacture of L-ascorbic acid from L-gulono-gamma-lactone.

EP 0 911 415 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 12 1547

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	WO 85 01745 A (KRAFT, INC.) 25 April 1985 * page 4, line 9 - line 25 * * page 5, line 5 - line 18 * * page 13, line 31 - page 14, line 20 * ---	1-7	C12P17/04
A	EP 0 207 763 A (BIO-TECHNICAL RESOURCES INC.) 7 January 1987 * page 2, line 19 - page 3, line 14 * * page 8, line 15 - line 23 * ---	1-7	
D,A	MORIMITSU NISHIKIMI ET AL.: "Occurrence in yeast of L-Galactonolactone oxidase which is similar to a key enzyme for ascorbic acid biosynthesis in animals, L-Gulonolactone Oxidase" ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, vol. 191, no. 1, December 1978, pages 479-486, XP002095259 * the whole document * -----	1-7	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C12P C12N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 2 March 1999	Examiner MONTERO LOPEZ B.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03 82 (P4/C01)